

ON SUBGRAPHS OF TRIPARTITE GRAPHS

ABHIJEET BHALKIKAR AND YI ZHAO

ABSTRACT. Bollobás, Erdős, and Szemerédi [Discrete Math 13 (1975), 97–107] investigated a tripartite generalization of the Zarankiewicz problem: what minimum degree forces a tripartite graph with n vertices in each part to contain an octahedral graph $K_3(2)$? They proved that $n + 2^{-1/2}n^{3/4}$ suffices and suggested it could be weakened to $n + cn^{1/2}$ for some constant $c > 0$. In this note we show that their method only gives $n + (1 + o(1))n^{11/12}$ and provide many constructions that show if true, $n + cn^{1/2}$ is better possible.

1. INTRODUCTION

Let K_t denote the complete graph on t vertices. As a foundation stone of extremal graph theory, Turán's theorem in 1941 [10] determines the maximum number of edges in graphs of a given order not containing K_t as a subgraph (the $t = 3$ case was proven by Mantel in 1907 [6]). In 1975 Bollobás, Erdős, and Szemerédi [1] investigated the following Turán-type problem for multipartite graphs.

Problem 1. *Given integers n and $3 \leq t \leq r$, what is the largest minimum degree $\delta(G)$ among all r -partite graphs G with parts of size n and which do not contain a copy of K_t ?*

The $r = t$ case of Problem 1 had been a central topic in Combinatorics until it was finally settled by Haxell and Szabó [4], and Szabó and Tardos [8]. Recently Lo, Treglown, and Zhao [9] solved many $r > t$ cases of the problem, including when $r \equiv -1 \pmod{t-1}$ and $r = \Omega(t^2)$.

For simplicity, let $G_r(n)$ denote an (arbitrary) r -partite graph with parts of size n . Let $K_r(s)$ denote the complete r -partite graph with parts of size s . In particular, $K_3(2)$ is known as the *Octahedral graph*. In the same paper Bollobás, Erdős, and Szemerédi [1] also asked the following question.

Problem 2. *Given a tripartite graph $G = G_3(n)$, what $\delta(G)$ guarantees a copy of $K_3(2)$?*

Problem 2 is a natural generalization of the well-known Zarankiewicz problem [11], whose symmetric version asks for the largest number of edges in a bipartite graph $G_2(n)$ that contains no $K_2(s)$ as a subgraph (in other words, $K_2(s)$ -free).

In [1, Corollary 2.7] the authors stated that $\delta(G) \geq n + 2^{-1/2}n^{3/4}$ guarantees a copy of $K_3(2)$. This follows from [1, Theorem 2.6], which handles the general case of $K_3(s)$ for arbitrary s . Unfortunately, there is a miscalculation in the proof of [1, Theorem 2.6] and thus the bound $\delta(G) \geq n + 2^{-1/2}n^{3/4}$ is unverified. We follow the approach of [1, Theorem 2.6] and obtain the following result.

Theorem 3. *Given an integer $s \geq 2$ and $\varepsilon > 0$, let n be sufficiently large. If $G = G_3(n)$ satisfies $\delta(G) \geq n + (1 + \varepsilon)(s - 1)^{1/(3s^2)}n^{1-1/(3s^2)}$, then G contains a copy of $K_3(s)$.*

In particular, Theorem 3 implies that every $G = G_3(n)$ with $\delta(G) \geq n + (1 + o(1))n^{11/12}$ contains a copy of $K_3(2)$. Using a result of Erdős on hypergraphs [3], we give a different proof of Theorem 3 under a slightly stronger condition $\delta(G) \geq n + (3n)^{1-1/(3s^2)}$. Thus $cn^{11/12}$ is a natural additive term for Problem 2 under typical approaches for extremal problems.

On the other hand, the authors of [1] conjectured that $\delta(G) \geq n + cn^{1/2}$ suffices for Problem 2. Although not explained in [1], they probably thought of Construction 10, a natural construction based on the one for the Zarankiewicz problem. We indeed find many non-isomorphic constructions, Construction 11, with the same minimum degree.

Proposition 4. *For any $n = q^2 + q + 1$ where q is a prime power, there are many tripartite graphs $G = G_3(n)$ such that $\delta(G) \geq n + n^{1/2}$ and G contains no $K_3(2)$.*

Key words and phrases. Turn-type problems, tripartite graphs, Zarankiewicz problem.

This paper is a result of the Undergraduate Research Initiations in Mathematics, Mathematics Education and Statistics (RIMMES) that the first author participated in under the guidance of the second author. The second author is partially supported by a Simons Collaboration Grant 710094.

Theorem 3 and Proposition 4 together show that the answer for Problem 2 lies between $n + n^{1/2}$ and $n + n^{11/12}$. The truth may be closer to the lower bound. If this is the case, then verifying it may be hard given the presence of many non-isomorphic constructions.

We know less about the minimum degree of $G_3(n)$ that forces a copy of $K_3(s)$. Theorem 3 shows that $\delta(G_3(n)) \geq n + cn^{1-1/(3s^2)}$ suffices. As shown in Remark 12, if there is a $K_2(s)$ -free bipartite graph $B = G_2(n)$ with $\delta(B) = \Omega(n^{1-1/s})$, then our constructions for Proposition 4 provide a tripartite $K_3(s)$ -free graph $G = G_3(n)$ with $\delta(G) = n + \Omega(n^{1-1/s})$.

2. PROOF OF THEOREM 3

In order to prove Theorem 3, we need the following results from [1].

Lemma 5. [1, Theorem 2.3] *Suppose every vertex of $G = G_3(n)$ has degree at least $n+t$ for some integer $t \leq n$. Then there are at least t^3 triangles in G .*

Lemma 6. [1, Lemma 2.4] *Let $X = \{1, \dots, N\}$ and $Y = \{1, \dots, p\}$. Suppose A_1, \dots, A_p are subsets of X such that $\sum_{i=1}^p |A_i| \geq pwN$ and $(1-\alpha)wp \geq q$, where $0 < \alpha < 1$ and N, p and q are natural numbers. Then there are q subsets A_{i_1}, \dots, A_{i_q} such that $|\bigcap_{j=1}^q A_{i_j}| \geq N(\alpha w)^q$.*

Let $z(n, s)$ denote the largest number of edges in a bipartite $K_2(s)$ -free graph with n vertices in each part. Kővári, Sós, and Turán [5] gave the following upper bound for $z(n, s)$.¹

Lemma 7. $z(n, s) \leq (s-1)^{1/s}(n-s+1)n^{1-1/s} + (s-1)n$.

We are ready to prove Theorem 3.

Proof of Theorem 3. Let G be a tripartite graph with three parts V_1, V_2, V_3 of size n each. Suppose $\delta(G) \geq n+t$, where $t = (1+\varepsilon)(s-1)^{\frac{1}{3s^2}}n^{1-\frac{1}{3s^2}} > n^{1-\frac{1}{3s^2}}$. By Lemma 5, G contains at least t^3 triangles.

We apply Lemma 6 in the following setting. Let $Y = V_1 = \{1, \dots, n\}$ and $X = V_2 \times V_3$ be the set of n^2 pairs (x, y) , $x \in V_2, y \in V_3$. For $1 \leq i \leq n$, let A_i be the set of pairs $(x, y) \in X$ for which $\{i, x, y\}$ spans a triangle of G . Then $\sum_{i=1}^n |A_i|$ is the number of triangles in G so $\sum_{i=1}^n |A_i| \geq t^3$. Let $N = n^2$, $p = n$, $q = s$, $w = t^3/n^3$, and $\alpha = 1/(1+\varepsilon)$. The assumptions of Lemma 6 hold because $pwN = t^3$ and

$$(1-\alpha)wp = \frac{\varepsilon}{1+\varepsilon} \left(\frac{t}{n}\right)^3 n > \frac{\varepsilon}{1+\varepsilon} n^{-1/s^2} n > s$$

as n is sufficiently large. By Lemma 6, there are $i_1, \dots, i_s \in V_1$ such that

$$\left| \bigcap_{j=1}^s A_{i_j} \right| \geq N(\alpha w)^q = n^2 \left(\frac{t^3}{(1+\varepsilon)n^3} \right)^s$$

Since

$$t > (1+\varepsilon)^{\frac{2}{3}}(s-1)^{\frac{1}{3s^2}}n^{1-\frac{1}{3s^2}} \quad \text{and} \quad \frac{t^3}{(1+\varepsilon)n^3} > (1+\varepsilon)(s-1)^{1/s^2}n^{-1/s^2},$$

we have

$$(1) \quad \left| \bigcap_{j=1}^s A_{i_j} \right| > (1+\varepsilon)^s (s-1)^{1/s} n^{2-1/s} \geq (s-1)^{1/s} n^{2-1/s} + (s-1)n.$$

Let B denote the bipartite graph between V_2 and V_3 with $E(B) = \bigcap_{j=1}^s A_{i_j}$. By (1) and Lemma 7, B contains a copy of $K_2(s)$. Since every edge of B forms a triangle with each of $i_1, \dots, i_s \in V_1$, this copy of $K_2(s)$ together with i_1, \dots, i_s span a desired copy of $K_3(s)$ in G . \square

We now give another proof of Theorem 3 with slightly larger $\delta(G)$ by a classical result of Erdős on hypergraphs [3]. An r -uniform hypergraph or r -graph is a hypergraph such that all its edges contain exactly r vertices. Let $K_r^r(s)$ denote the complete r -partite r -graph with s vertices in each part, namely, its vertex set consists of disjoint parts V_1, \dots, V_r of size s , and edges set consists of all r -sets $\{v_1, \dots, v_r\}$ with $v_i \in V_i$ for all i .

Lemma 8. [3, Theorem 1] *Given integers $r, s \geq 2$, let n be sufficiently large. Then every r -graph on n vertices with at least $n^{r-s^{1-r}}$ edges contains a copy of $K_r^r(s)$.*

¹In [1] the authors instead used the Turán number $\text{ex}(2n, K_2(s))$, which gives a slightly worse constant here.

Proposition 9. *Let $s \geq 2$ and n be sufficiently large. Every tripartite graph $G = G_3(n)$ with $\delta(G) \geq n + (3n)^{1-1/(3s^2)}$ contains a copy of $K_3(s)$.*

Proof. Suppose $G = G_3(n)$ satisfies $\delta(G) \geq n + (3n)^{1-1/(3s^2)}$. By Lemma 5, G contains at least $(3n)^{3-1/s^2}$ triangles. Let H be the 3-graph on $V(G)$, whose edges are triangles of G . Then H has $3n$ vertices and at least $(3n)^{3-s^{-2}}$ edges. By Lemma 8 with $r = 3$ and $s = 2$, H contains a copy of $K_3^3(s)$, which gives a copy of $K_3(s)$ in G . \square

3. PROOF OF PROPOSITION 4

In this section we prove Proposition 4 by constructing many tripartite $K_3(2)$ -free graphs $G_3(n)$ with $\delta(G_3(n)) \geq n + n^{1/2}$.

One main building block is a bipartite $K_2(2)$ -free graph $G_0 = G_2(n)$ with $\delta(G_0) \geq \sqrt{n}$. First shown in [7], such a graph exists when $n = q^2 + q + 1$ and a projective plane of order q exists. Indeed, two parts of $V(G)$ correspond to the points and lines of the projective plane and a point is adjacent to a line if and only if the point lies on the line. It is easy to see that such graph contains no $K_2(2)$ and is regular with degree $q + 1 > \sqrt{n}$.

Construction 10. *Suppose $G = G_3(n)$ has parts V_1, V_2 and V_3 each of size n . Let the bipartite graphs between V_1 and V_2 and between V_1 and V_3 be complete, while the bipartite graph between V_2 and V_3 is G_0 defined above.*

Since $\deg_{G_0}(v) \geq \sqrt{n}$ for $v \in V_2 \cup V_3$, we have $\delta(G) \geq n + \sqrt{n}$. Furthermore, G contains no $K_3(2)$ because by the definition of G_0 , there is no $K_2(2)$ between V_2 and V_3 .

We now provide a family of constructions with the same properties.




FIGURE 1. Graph from Construction 11

Construction 11. *Let $G = G_3(n)$ be a tripartite graph with parts V_1, V_2 , and V_3 of size n each. Partition $V_2 = X_2 \cup Y_2$ arbitrarily such that $\sqrt{n} \leq |X_2| \leq |Y_2|$. Partition $V_3 = X_3 \cup Y_3$ arbitrarily such that $|X_3| = |Y_2|$ and $|Y_3| = |X_2|$.*

The bipartite graphs (V_1, X_2) , (X_2, Y_3) , (Y_3, Y_2) , (Y_2, X_3) , and (X_3, V_1) are complete, in other words, V_1, X_2, Y_3, Y_2, X_3 form a blowup of C_5 . Let the bipartite graph between V_1 and $Y_2 \cup Y_3$ be isomorphic to G_0 (note that $|X_2| + |Y_2| = |X_3| + |Y_3| = n$).

For any vertex $v \in X_2$, $\deg(v) = |V_1| + |Y_3| \geq n + \sqrt{n}$. The vertices $v \in X_3$ satisfy $\deg(v) = |V_1| + |Y_2| \geq n + n/2$. For any $v \in Y_2$, $\deg(v) \geq |V_3| + \delta(G_0) \geq n + \sqrt{n}$. The same holds for the vertices of Y_3 . At last, every vertex $v \in V_1$ satisfies $\deg(v) \geq |X_2| + |X_3| + \delta(G_0) \geq n + \sqrt{n}$. These together show that $\delta(G) \geq n + \sqrt{n}$.

Suppose G contains a copy of $K_3(2)$ with vertex set S . Then $|S \cap V_i| = 2$ for $i = 1, 2, 3$. Since there is no edge between X_2 and X_3 , either $S \cap X_2 = \emptyset$ or $S \cap X_3 = \emptyset$. Suppose, say, $S \cap X_2 = \emptyset$, which forces $|S \cap Y_2| = 2$. Hence $S \cap Y_2$ and $S \cap V_1$ span a copy of $K_2(2)$, contradicting the definition of G_0 .

If letting $X_2 = \emptyset = Y_3$ in Construction 11, then we obtain Construction 10. Nevertheless, we prefer viewing Constructions 10 and 11 as different constructions because after removing $o(n^2)$ edges, Construction 11 contains many 5-cycles while Construction 10 does not.

Remark 12. *If we replace G_0 by a $K_2(s)$ -free bipartite graph with n vertices in each part in Constructions 10 and 11, then we obtain a $K_3(s)$ -free tripartite graph $G_3(n)$. It has been conjectured that there exist a $K_2(s)$ -free bipartite graph with n vertices in each part and $\Omega(n^{2-1/s})$ edges (this is known for $s = 2, 3$ [2, 7]). If such bipartite graph exists and is regular, then (revised) Constructions 10 and 11 provide a $K_3(s)$ -free tripartite graph $G = G_3(n)$ with $\delta(G) = n + \Omega(n^{1-1/s})$.*

REFERENCES

- [1] B. Bollobás, P. Erdős, A. Szemerédi. On complete subgraphs of r -chromatic graphs. *Discrete Math*, 13:97-107, 1975.
- [2] W. G. Brown. On graphs that do not contain a Thomsen graph. *Canada Math. Bull.* 9, 281285, 1966.
- [3] P. Erdős. On extremal problems of graphs and generalized graphs. *Israel Journal of Mathematics*, 2, 3:183-190, 1964.
- [4] P. Haxell and T. Szabó. Odd independent transversals are odd. *Combin. Probab. Comput.*, 15(1-2):193–211, 2006.
- [5] T. Kővári, V. Sós, P. Turán. On a problem of K. Zarankiewicz, *Colloquium Math.*, 3:5057, 1954.
- [6] W. Mantel. Problem 28 (Solution by H. Gouwentak, W. Mantel, J. Teixeira de Mattes, F. Schuh and W. A. Wythoff), *Wiskundige Opgaven*, 10:6061, 1907.
- [7] I. Rieman. Über ein Problem von K. Zarankiewicz. *Acta Math. Acad. Sci. Hungar.* 9, 3-4:269273, 1958.
- [8] T. Szabó and G. Tardos. Extremal problems for transversals in graphs with bounded degree. *Combinatorica*, 26(3):333–351, 2006.
- [9] A. Lo, A. Treglown and Y. Zhao. Complete subgraphs in a multipartite graph *Combin. Probab. Comput.*, to appear.
- [10] P. Turán. On an extremal problem in graph theory. *Mat. Fiz. Lapok*, 48:436452, 1941.
- [11] K. Zarankiewicz. Problem P 101. *Colloq. Math.*, 2:301, 1951.

DEPARTMENT OF MATHEMATICS AND STATISTICS, GEORGIA STATE UNIVERSITY, ATLANTA, GA 30303
E-mail address: `abhalkikar1@student.gsu.edu`

DEPARTMENT OF MATHEMATICS AND STATISTICS, GEORGIA STATE UNIVERSITY, ATLANTA, GA 30303
E-mail address: `yzhao6@gsu.edu`